

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 August 2002 (22.08.2002)

PCT

(10) International Publication Number
WO 02/064183 A1

(51) International Patent Classification⁷: **A61L 31/08**,
G02B 1/04

(21) International Application Number: PCT/US01/04524

(22) International Filing Date: 9 February 2001 (09.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(71) Applicant: **JOHNSON & JOHNSON VISION CARE, INC.** [US/US]; 7500 Centurion Parkway, Suite 100, Jacksonville, FL 32256 (US).

(72) Inventors: **LAN, Jenny**; 1/227 Rawson St., Auburn, New South Wales 2144 (AU). **VANDERLAAN, Doug**; 8114 Parkridge Circle S., Jacksonville, FL 32211 (US). **WILLCOX, Mark**; 22 College St, Balmain, New South Wales 2041 (AU). **ALIWARGA, Yulina**; 81 Ruthven St, Bondi Junction, New South Wales 2022 (AU).

(74) Agents: **JOHNSON, Philip, S. et al.**; Johnson & Johnson, One Johnson & Johnson Plaza, New Brunswick, NJ 08933 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: BIOMEDICAL DEVICES WITH ANTIMICROBIAL CATIONIC PEPTIDE AND PROTEIN COATINGS

(57) Abstract: Biomedical devices with antimicrobial coatings are provided. One or more surfaces of the device are coated with a cationic peptide, cationic proteins, or mixtures thereof to impart antimicrobial properties to the surface.

WO 02/064183 A1

BIOMEDICAL DEVICES WITH ANTIMICROBIAL CATIONIC PEPTIDE COATINGS

5

Field of the Invention

This invention relates to coated devices. In particular, the invention provides biomedical devices on the surfaces of which antimicrobial coatings of a cationic peptide, a cationic protein, or both are formed.

10

Background of the Invention

Devices for use in and on the human body are well known. The chemical composition of the surfaces of such devices plays a pivotal role in dictating the overall efficacy of the devices. Additionally, it is known that providing such devices with an antimicrobial surface is advantageous.

15

A wide variety of bactericidal and bacteriostatic coatings have been developed. For example, cationic antibiotics, such as polymyxin, vancomycin, and tetracycline have been used as coatings for contact lenses. Further, metal chelating agents, substituted and unsubstituted polyhydric phenols, aminophenols, alcohols, acid and amine derivatives, and quarternary ammonium have been used as antimicrobial agents for contact lenses.

20

However, the use of these known antimicrobial coatings has disadvantages. With the use of antibiotic coatings, microorganisms resistant to the antibiotics may develop. Chelating agent use fails to address the numbers of bacteria that adhere to the device. Some of the prior art coatings, for example phenol derivatives and cresols, can produce ocular toxicity or allergic reactions. Quarternary ammonium compounds are problematic because of their irritancy. Thus, a need exists for safe and effective antimicrobial coatings that overcomes at least some of these disadvantages.

25
30

Detailed Description of the Invention and Preferred Embodiments

The present invention provides biomedical devices with an antimicrobial coating and processes for the production of the biomedical devices. It is an
5 unexpected discovery of the invention that certain cationic peptides, cationic proteins, or both may be used to provide antimicrobial coatings for biomedical devices. In particular, it is one discovery of the invention that protamine, melittin, cecropin A, nisin, or combinations thereof, when used as surface coatings, reduce adherence of bacteria to a device's surface, reduce growth of bacteria adhered to a
10 device, or both by greater than about 50 percent.

In one embodiment, the invention provides a biomedical device comprising, consisting essentially of, and consisting of at least one surface comprising, consisting essentially of, and consisting of a coating effective amount of one of protamine,
15 melittin, cecropin A, nisin, or combinations thereof. In yet another embodiment, a method for manufacturing biomedical devices comprising, consisting essentially of, and consisting of contacting at least one surface of a biomedical device with a coating effective amount of protamine, melittin, cecropin A, nisin, or combinations thereof is provided.

20

By "biomedical device" is meant any device designed to be used while in or on either or both human tissue or fluid. Examples of such devices include, without limitation, stents, implants, catheters, and ophthalmic lenses. In a preferred embodiment, the biomedical device is an ophthalmic lens including, without
25 limitation, contact or intraocular lenses. More preferably, the device is a contact lens, most preferably a soft contact lens.

Protamine is isolatable from the sperm of a variety of animals including, without limitation, man. Melittin is isolatable from bee venom. Cecropin A and
30 nisin are isolatable from *Aedes aegypti* and *Lactococcus lactis*, respectively. All four

3

are members of a broad group of cationic peptides and proteins which group includes, without limitation, defensins, magainins, and colicins. It is an unexpected discovery of this invention that only certain cationic peptides and proteins significantly reduce bacterial adherence, bacterial growth, or both on biomedical
5 devices.

Protamine, melittin, cecropin A, and nisin useful in the invention are all commercially available. Alternatively, these cationic peptides and proteins may be produced by known means. For purposes of the invention, generally the purity of the
10 cationic peptide used is at least about 75 %, preferably at least about 90 %.

Protamine, melittin, cecropin A, nisin, or combinations thereof may be adsorbed to polymer surfaces of a biomedical device. The cationic peptides and proteins may be used on any surface, but most advantageously are used with
15 negatively charged surfaces.

The cationic peptides and proteins alternatively may be bound to the polymer surfaces. This may be either a direct reaction or, preferably, a reaction in which a coupling agent is used. For example, a direct reaction may be accomplished by the
20 use of a reagent of reaction that activates a group in the surface polymer or the cationic peptide making it reactive with a functional group on the peptide or polymer, respectively, without the incorporation of a coupling agent. For example, one or more amine groups on the peptide may be reacted directly with isothiocyanate, acyl azide, N-hydroxysuccinimide ester, sulfonyl chloride, an aldehyde, glyoxal epoxide,
25 carbonate, aryl halide, imido ester, or an anhydride group on the polymer.

In an alternative embodiment, coupling agents may be used. Coupling agents useful for coupling the cationic peptide or protein to the device's surface include, without limitation, N, N'-carbonyldiimidazole, carbodiimides such as 1-ethyl-3-(3-
30 dimethylaminopropyl)carbodiimide ("EDC"), dicyclohexyl carbodiimide, 1-

cyclohexyl-3-(2-morpholinoethyl)carbodiimide, diisopropyl carbodiimide, or mixtures thereof. The carbodiimides also may be used with N-hydroxysuccinimide or N-hydroxysulfosuccinimide to form esters that can react with amines to form amides.

5 Amino groups also may be coupled to the polymer by the formation of Schiff bases that can be reduced with agents such as sodium cyanoborohydride and the like to form hydrolytically stable amine links. Coupling agents useful for this purpose include, without limitation, N-hydroxysuccinimide esters, such as dithiobis(succinimidylpropionate), 3,3'-dithiobis(sulfosuccinimidylpropionate),
10 disuccinimidyl suberate, bis(sulfosuccinimidyl) suberate, disuccinimidyl tartarate and the like, imidoesters, including, without limitation, dimethyl adipimate, difluorobenzene derivatives, including without limitation 1,5-difluoro-2,4-dinitrobenzene, bromofunctional aldehydes, including without limitation gluteraldehyde, and bis epoxides, including without limitation 1,4-butanediol
15 diglycidyl ether. One ordinarily skilled in the art will recognize that any number of other coupling agents may be used depending on the functional groups present on the device's surface.

One ordinarily skilled in the art also will recognize that, if the device's surface
20 does not contain suitable reactive groups, such suitable groups may be incorporated into the polymer by any conventional organic synthesis methods. Alternatively, the reactive groups may be introduced by the addition of polymerizable monomers containing reactive groups into the monomer mixture used to form the polymer.

25 Examples of polymer surfaces onto which the cationic peptides and proteins may be adsorbed or bonded are surfaces formed from, without limitation, polymers and copolymers of styrene and substituted styrenes, ethylene, propylene, acrylates and methacrylates, N-vinyl lactams, acrylamides and methacrylamides, acrylonitrile, acrylic and methacrylic acids as well as polyurethanes, polyesters,
30 polydimethylsiloxanes and mixtures thereof. Such polymers may include hydrogels

and silicone containing hydrogels. Preferably, lightly crosslinked polymers and copolymers of 2-hydroxyethylmethacrylate ("HEMA") are used. By "lightly crosslinked" is meant that the polymer has a low enough crosslink density so that it is soft and elastic at room temperature. Typically, a lightly crosslinked polymer will have about 0.1 to about 1 crosslinking molecule per about 100 repeating monomer units. Examples of suitable lightly crosslinked HEMA polymers and copolymers include without limitation, etafilcon A and copolymers of glycerol methacrylate and HEMA. Also preferably, silicone hydrogels, especially those of hydrophilic monomers, such as N,N-dimethylacrylamide, are used.

10

In one embodiment of the process for making the device of the invention, the surface to be coated is contacted with the protamine, melittin, cecropin A, nisin or combinations thereof in any convenient manner. Preferably, mixtures of protamine and melittin are used. For example, the device may be placed in a solution of protamine and solvent into which the medical device is placed. As an alternative, the device's surface may first be treated with a coupling agent and the surface then placed in a solution of the selected cationic peptide or protein. As yet another alternative the peptide or protein may be reacted alone with the polymer surface.

20

Suitable solvents for use in the invention are those that are capable of dissolving protamine, melittin, cecropin A, or nisin singly or in combination. Preferably, the coating process is carried out in water, alcohol, or mixtures thereof. EDC is effective in aqueous solutions and, thus, is a preferred coupling agent.

25

The coupling agents may be used alone or in combination with agents capable of stabilizing any reactive intermediate formed. For example, EDC may be used with N-hydroxysuccinimide as a stabilizer. Additionally, it may be desirable to adjust the pH. Preferably, the pH is adjusted to about 6.5 to about 8.0, more preferably about 7.0 to about 7.5.

30

A coupling effective amount of a coupling agent is used which amount is an amount sufficient to couple the peptide or protein to the device surface. The precise amount of coupling agent used will depend on the surface's chemistry as well as the agent selected. Generally, about 0.01 to about 10 weight percent, preferably about 5 0.01 to about 5.0, more preferably about 0.01 to about 1 weight percent of the coupling agent is used based on the weight of the coating solution. By coating solution is meant the peptide or protein with one or more of the solvent, coupling agent, buffer, and the like. Typically, the amount of coating solution used per lens will be about 1 ml to about 10 ml, preferably about 1 ml to about 5 ml, more 10 preferably about 1 ml to about 2 ml per lens.

In the processes of the invention, a coating effective amount of protamine, melittin, cecropin A, nisin, or combinations thereof is used meaning an amount that when contacted with the surface is sufficient to coat the surface so as to impart the 15 desired antimicrobial properties to the surface. By antimicrobial properties is meant either or both the ability to significantly reduce, meaning by greater than about 50 percent, either or both the amount of bacteria adhering to the surface and the growth of bacteria adhered to the surface. In the case of contact lenses, generally, the amount contacted with the lens is about 1 μ g to about 10 mg, preferably about 10 μ g 20 to about 1 mg per lens. The amount of coating resulting per contact lens is about 50 to about 1000 μ g. In cases in which combinations of melittin and protamine are used, the amount of protamine used preferably is about 500 μ g/ml or less.

Temperature and pressure are not critical to the processes of the invention 25 and the process may be conveniently carried out at room temperature and pressure. The contact time used will be a length of time sufficient to coat the surface to the extent desired. Preferably, contact time is about 60 seconds to about 24 hours.

Following contacting, the surface may be washed with water or buffered 30 saline solution to remove unreacted protamine, melittin, colicin and solvent. One

ordinarily skilled in the art will recognize that the polymer for producing the surface to be coated by the method of the invention may contain other monomers and additives. For example, ultra-violet absorbing monomers, reactive tints, processing aids, and the like may be used.

5

The invention will be further clarified by a consideration of the following, non-limiting examples.

Examples

In the following examples, the cationic proteins and peptides listed on Table 1 were used.

10

Table 1

CATIONIC PEPTIDE/PROTEIN	SOURCE
Protamine	Fish
Cecropin A	Insect
Cecropin P1	Pig
Melittin	Insect
Melittin-1-13 AA	Synthetic
Magainin 1	Frog
Magainin 2	Frog
Defensin HNP-1	Human
β Defensin 1	Human
Secretory leukocyte protease inhibitor (SLPI)	Human
Colicin	Gram - bacteria
Nisin	Gram + bacteria

15

Bacterial strains used in the Examples are listed on Table 2.

Table 2

Strain	Isolation Site
<i>Pseudomonas aeruginosa</i> Paer1	CLARE*
<i>Pseudomonas aeruginosa</i> 6294	MK ⁺ -ULCER
<i>Pseudomonas aeruginosa</i> 6206	MK-ulcer
<i>Pseudomonas aeruginosa</i> ATCC 15442	Environmental strain
<i>Serratia marcescens</i> Smar5	CLARE**
<i>Escherichia coli</i> Ecol8	CLARE
<i>Staphylococcus intermedius</i> Sint 2	Asymptomatic lens
<i>Staphylococcus aureus</i> Saur31	CLPU***

*By "MK" is meant microbial keratitis.

**By "CLARE" is meant contact lens induced red eye.

*** By "CLPU" is meant contact lens-induced peripheral ulcers.

20

The majority of testing was performed with strains *P. aeruginosa* 6294 and *S. aureus* 31. *P. aeruginosa* and *S. aureus* are the most common bacteria causing eye inflammation or infections for contact lens wearers. Other strains were used to validate results or assess the effectiveness of the compounds over a range of bacteria.

Example 1

To assess the effect of the cationic proteins/peptides in solution against rapid growing bacterial cells, bacteria were cultured in Trypticase soy broth ("TSB") overnight at 35°C. Aliquots (20µl) of this cell suspension were then added to fresh TSB (10ml). Different concentrations of the cationic proteins/peptides were added to the fresh broth and incubated for up to 48h at 35°C. Samples were taken at different time points and the optical density at 660nm measured as a measure of changes in bacterial numbers was measured.

15

To assess the effect of the cationic proteins/peptides in solution against slow growing bacterial cells, cells were grown as previously in TSB. The cells were then harvested by centrifugation and washed in phosphate buffered saline (PBS; NaCl 8g/l; KCl 0.2g/l; Na₂HPO₄ 1.15g/l; KH₂PO₄ 0.2g/l). The cells were then re-suspended to OD 0.1 (unless otherwise stated) at 660nm in PBS, different concentrations of cationic proteins/peptides were added and incubated for up to 48h at 35°C. Samples were taken at different points and numbers of bacteria analyzed using the Miles and Misra technique (i.e. numbers that are viable after plating dilutions onto nutrient agar plates).

25

The results obtained in solution for all cationic proteins and peptides studied are shown on Table 3. Most protein/peptides were screened against only *P. aeruginosa* 6294 and *S. aureus* 31. If these showed reductions in growth then other strains may have been examined. As can be seen from Table 3, protamine and melittin were the most efficacious at preventing the growth of gram-positive and gram-negative bacteria in solution.

30

In Table 3, slow growing cells are those re-suspended in PBS with cationic protein/peptide and rapid growing are those re-suspended in TSB plus cationic protein/peptide. The numbers in $\mu\text{g/ml}$ are in concentration showing peak activity.

- 5 A “-” sign indicates no reduction in bacterial growth, a “+” sign indicates a 1 to 50 % reduction in growth, a “++” sign indicates a 51 to 89 % reduction in growth, a “+++” sign indicates a 90 to 98 % reduction in growth a “++++” sign indicates a greater than 98 % reduction in growth.

10

15

20

25

Table 3

	Paer 1	6294	6206	Saur 31	Sint 2	Smar 5	Ecol 8
Protamine 125- 1000µg/ml	++++ (slow growing) - (rapid growing)	++++ (slow growing)	++++ (slow growing)	++++ (slow growing)	++++ (slow growing) - (rapid growing)	++++ (1000 µg/ml slow growing) - (<1000 µg/ml slow growing)	ND
Melittin 15µg/ml	++++ (24h, 15µg/ml slow growing) - (48h rapid growing)	++ (24h, 15µg/ml slow growing) - (48h slow growing)	ND	++++ (15µg/ml rapid growing) ++++ (24h, 15µg/ml slow growing)	++++ (24h, 15µg/ml slow growing) ++++ (24h, 15µg/ml rapid growing)	-	ND
Magainin 1 10- 200µg/ml	-	++ (slow growing)	ND	-	-	ND	ND
Magainin 2 100- 400µg/ml	ND	+++ (24h, 400µg/ml, slow growing) - (48h, 400µg/ml, rapid growing)	ND	-	ND	ND	ND
Cecropin P1 1- 50µg/ml	+ (slow growing) - (rapid growing)	-	ND	-/+ (rapid growing) - (slow growing)	-	ND	ND
Cecropin A 4- 60µg/ml	ND	++++ (slow growing)	ND	-	ND	ND	ND
SLPI 10- 100µg/ml	-	-	-	-	-	ND	ND
Colicin 1- 10units/ml	ND	++ (24h)	ND	++ (24h)	ND	ND	- (24h)
Nisin 32µg/ml	ND	++ (2µg/ml)	ND	-	ND	ND	ND
β Defensin 60µg/ml	ND	++ (24h, 60µg/ml)	ND	-	ND	ND	ND
Melittin 1 13- 1000µg/ml	ND	++ (24h, 1000µg/ml)	ND	++ (48h, 24h, 1000µg/ml)	ND	ND	ND
Defensin HNP1 1- 10µg/ml	ND	+ (2µg/ml)	ND	-	ND	ND	ND
Transferrin 125- 2000µg/ml	ND	++	ND	-	ND	ND	ND

* By "ND" is meant not determined.

Example 2

For conducting total counts, etafilcon A lenses were removed from the
5 manufacturers vials, washed three times in 1ml PBS and then coated with various
concentrations of cationic proteins/peptides overnight at 37°C either individually or in
combination. After incubation, the lenses were washed three times in PBS and 0.5ml
of 1×10^8 bacterial cells/ml was added to the lenses. After incubation at ambient
temperature for 10min, the lenses were washed three times in PBS to remove non-
10 adherent or loosely adherent bacteria and stained with crystal violet for 1 min. The
number of cells per lens was examined under the microscope. Five grids (0.005625
 mm^2) per lens were counted and triplicate lenses for each treatment were assayed.

For conducting viable counts, etafilcon A lenses were removed from the
15 manufacturers vials, washed three times in 1ml PBS and then coated with various
concentrations of cationic proteins/peptides overnight at 37°C (either individually or
in combination). After incubation, the lenses were washed three times in PBS and
0.5ml of 1×10^8 bacterial cells/ml was added to the lenses. After incubation at ambient
temperature for 10min, the lenses were washed three times in PBS to remove non-
20 adherent or loosely adherent bacteria. Lenses were then homogenized using 1 ml PBS
and a small magnetic stirring bar (octagonal cross-section, 0.5" X 0.125") and stirred at
maximum speed for one hour which was sufficient for lens disintegration. Serial dilutions
were then made according to the technique of Miles and Misra and aliquots (20 μL)
plated out on nutrient agar. After incubation overnight at 37°C, viable bacteria were
25 determined and results expressed as colony forming units/ mm^2 after calculation of the
surface area of the lens (approximately 310mm^2).

The lenses were incubated in concentrations of cationic proteins/peptides that
were either effective in solution or the highest concentration available if there was no
30 effect in solution. After rinsing, bacteria were added and numbers of cells analyzed as the

12

total cells per mm² of the lens or the number of viable cells per mm² of the lens. The results are shown on Table 4.

Table 4

Peptide	Strain	Reduction v. control*	Reduction v total**
Protamine	6294	80%	80%
	Saur 31	-	-
Melittin	6294	70%	-
	Saur 31	60%	-
Magainin 1	6294	-	-
	Saur 31	-	-
Magainin 2	6294	-	-
	Saur 31	-	-
Cecropin P1	6294	-	-
	Saur 31	-	-
Cecropin A	6294	-	93%
	Saur 31	-	-
SLPI	6294	-	-
	Saur 31	-	-
Colicin	6294	-	-
	Saur 31	-	-
Nisin	Saur 31	-	50%
β Defensin 1	6294	-	-
	Saur 31	-	-
Melittin-1-13 AA	6294	-	-
	Saur 31	-	-
Defensin HNP-1	6294	-	-
	Saur 31	-	-
Protamine 1000 μ g/ml and Melittin 15 μ g/ml	6294	90%	-
	Saur 31	-	-
Protamine 500 μ g/ml and Melittin 15 μ g/ml	6294	65%	-
	Saur 31	60%	-
Transferrin (125 μ g/ml)	6294	-	-

5 *Effect compared to adhesion control lens that was not coated with cationic protein/peptide before bacterial adhesion.

**Effect compared to the number of total bacterial cells adhered to lens coated with cationic proteins/peptides, i.e. the cationic prevented the growth of the adhered bacteria even though there may have been an increase in total cell numbers.

“-“ indicates no reduction in adhesion.

10

The results on Table 4 show that although nisin and cecropin A did not reduce the total adhesion of bacteria by increasing the total number of cells on the lens, there was a significant reduction in the viability of those cells compared to the

cells adhered to the uncoated lens. This indicates that the adhered bacteria were prevented from growing. Protamine significantly reduced the adhesion of *P. aeruginosa* 6294 to the lenses and also reduced the viability of the cells on the coated lenses. Melittin both reduced initial adhesion of *S. aureus* 31. A similar effect was
5 seen for *P. aeruginosa* 6294 and when a mixture of protamine and melittin was used.

What is claimed is:

1. A device comprising a biomedical device at least one surface of which
5 comprises a coating effective amount of one of protamine, melittin, cecropin A, nisin, or a combination thereof.
2. The device of claim 1 wherein the biomedical device is a contact lens.
- 10 3. The device of claim 1, wherein the at least one surface comprises a coating effective amount of protamine.
4. The device of claim 1, wherein the at least one surface comprises a coating effective amount of melittin.
- 15 5. The device of claim 1, wherein the at least one surface comprises a coating effective amount of protamine and melittin.
6. The device of claim 1, whereon the at least one surface comprises a coating
20 effective amount of cecropin A.
7. The device of claim 1, whereon the at least one surface comprises a coating effective amount of nisin.
- 25 8. The device of claim 1, wherein the surface further comprises a polymer selected from the group consisting of hydrogels, silicone containing hydrogels, polymers and copolymers of 2-hydroxyethylmethacrylate and mixtures thereof
9. The device of claim 8 wherein the polymer is a hydrogel.

10. The device of claim 8 wherein the polymer is a silicone containing hydrogel.

11. The device of claim 8 wherein the polymer is a polymer or copolymer of 2-hydroxyethylmethacrylate.

5

12. The device of claim 11 wherein the copolymer of 2-hydroxyethylmethacrylate is a lightly crosslinked copolymer of 2-hydroxyethylmethacrylate.

13. A contact lens at least one surface of which comprises a coating effective
10 amount of protamine, melittin, cecropin A, nisin, or a combination thereof.

14. The contact lens of claim 13 wherein the surface further comprises a polymer selected from the group consisting of hydrogels, silicone containing hydrogels, polymers and copolymers of 2-hydroxyethylmethacrylate and mixtures thereof

15

15. The contact lens of claim 14 wherein the polymer is a hydrogel.

16. The contact lens of claim 14 wherein the polymer is a silicone containing hydrogel.

20

17. The contact lens of claim 14 wherein the polymer is a polymer or copolymer of 2-hydroxyethylmethacrylate.

18. The contact lens of claim 17 wherein the copolymer of 2-hydroxyethylmethacrylate is a lightly crosslinked copolymer of 2-hydroxyethylmethacrylate.

25

19. A process for manufacturing a device comprising the step of contacting at least one surface of a biomedical device with a coating effective amount of
30 protamine, melittin, cecropin A, nisin, or a combination thereof.

20. The process of claim 19 wherein the biomedical device is a contact lens.
21. The process of claim 20, further comprising the step of contacting the at least
5 one surface with a coupling effective amount of a coupling agent.
22. The process of claim 20, wherein the at least one surface is contacted with a
coating effective amount of protamine.
- 10 23. The process of claim 20, wherein the at least one surface is contacted with a
coating effective amount of melittin.
24. The process of claim 20, wherein the at least one surface is contacted with a
coating effective amount of protamine and melittin.
- 15 25. The process of claim 20, wherein the at least one surface is contacted with a
coating effective amount of cecropin A.
26. The process of claim 20, wherein the at least one surface is contacted with a
20 coating effective amount of nisin.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/04524

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 A61L31/08 G02B1/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61L C11D A61K G02B C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, COMPENDEX, CHEM ABS Data, EMBASE, MEDLINE, SCISEARCH

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 990 924 A (JOHNSON & JOHNSON VISION PROD) 5 April 2000 (2000-04-05) paragraphs '0006!-'0008! claims	1-3, 5, 8-21, 24
X	US 5 260 271 A (BLACKBURN PETER ET AL) 9 November 1993 (1993-11-09) column 3, line 15 - line 29 claim 7	1, 2, 7-21, 26
X	WO 98 40401 A (FRASER JANET R ;MCNICOL PATRICIA J (CA); MICROLOGIX BIOTECH INC (C) 17 September 1998 (1998-09-17) page 2, paragraphs 2,3 page 3, paragraph 2 table 1 claims 16-19	1, 2, 4, 6-21, 23, 25, 26
-/-		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

11 January 2002

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Muñoz, M

INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/US 01/04524

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 25183 A (ALLERGAN INC) 22 August 1996 (1996-08-22) page 7, line 12 - line 20 page 10, line 29 -page 11, line 7	1,2,6, 8-21,25
A	claims	4,23
X	US 4 704 131 A (NOISHIKI YASU HARU ET AL) 3 November 1987 (1987-11-03) claim 1	1,3,19, 22
A	WO 96 06532 A (NOVONORDISK AS ;JOHANSEN CHARLOTTE (DK)) 7 March 1996 (1996-03-07) page 2, paragraphs 2,3 claim 22	1-3,5, 13,19, 20,22,24
A	WO 98 40091 A (UNIV NEW YORK) 17 September 1998 (1998-09-17) example 8 claims 1,17,18 page 3, last paragraph -page 4, paragraph 1	1,13,19
A	US 5 786 324 A (GRAY BEULAH ET AL) 28 July 1998 (1998-07-28) column 10, line 46 - line 64 column 16, line 39 -column 17, line 31 claims 26,27	1,13,19
A	TEICHMAN JOEL M H ET AL: "Protamine sulfate and vancomycin are synergistic against staphylococcus epidermidis prosthesis infection in vivo." JOURNAL OF UROLOGY, vol. 152, no. 1, 1994, pages 213-216, XP001028063 ISSN: 0022-5347 abstract	1-6,13, 19,20,22
A	DATABASE WPI Section Ch, Week 199712 Derwent Publications Ltd., London, GB; Class D16, AN 1997-126705 XP002179130 & JP 09 010288 A (SUN CONTACT LENS KK), 14 January 1997 (1997-01-14) abstract	1-3,13, 19,20,22
T	ALIWARGA Y ET AL.: "Antimicrobial peptides: a potential role in ocular therapy" CLINICAL AND EXPERIMENTAL OPHTHALMOLOGY, vol. 29, no. 3, June 2001 (2001-06), pages 157-160, XP002179129 the whole document	1-3,5, 8-21,24

INTERNATIONAL SEARCH REPORT

national application No.
PCT/US 01/04524

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1,2,8-21(partial); 3,5,22,24 (complete)

A biomedical device or a contact lens comprising a coating of protamine

2. Claims: 1,2,8-21(partial); 4, 23 (complete)

A biomedical device or a contact lens comprising a coating of melittin

3. Claims: 1,2,8-21(partial); 6,25 (complete)

A biomedical device or a contact lens comprising a coating of cecropin A

4. Claims: 1,2,8-21(partial); 7,26 (complete)

A biomedical device or a contact lens comprising a coating of nisin

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int al Application No

PCT/US 01/04524

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0990924	A	05-04-2000	AU 5018599 A	06-04-2000
			BR 9904394 A	14-11-2000
			CN 1254729 A	31-05-2000
			EP 0990924 A1	05-04-2000
			JP 2000187187 A	04-07-2000
US 5260271	A	09-11-1993	CS 8906897 A2	16-07-1991
			US 5691301 A	25-11-1997
			US 5753614 A	19-05-1998
			AT 101490 T	15-03-1994
			AT 142504 T	15-09-1996
			AU 631803 B2	10-12-1992
			AU 3843089 A	12-01-1990
			DE 68913189 D1	24-03-1994
			DE 68913189 T2	19-05-1994
			DE 68927189 D1	17-10-1996
			DE 68927189 T2	30-01-1997
			DK 45690 A	21-02-1990
			EP 0382814 A1	22-08-1990
			EP 0545911 A2	09-06-1993
			FI 98880 B	30-05-1997
			HU 53795 A2	28-12-1990
			IE 63998 B1	28-06-1995
			IL 90700 A	24-06-1994
			JP 8009525 B	31-01-1996
			JP 3500051 T	10-01-1991
			NO 179354 B	17-06-1996
			NZ 229674 A	23-12-1992
			RU 2092180 C1	10-10-1997
			US 5304540 A	19-04-1994
			US 5334582 A	02-08-1994
			WO 8912399 A1	28-12-1989
			US 5135910 A	04-08-1992
			US 5217950 A	08-06-1993
			ZA 8904691 A	27-06-1990
			IE 940624 L	22-12-1989
WO 9840401	A	17-09-1998	US 6180604 B1	30-01-2001
			AU 6604798 A	29-09-1998
			WO 9840401 A2	17-09-1998
			EP 0966481 A2	29-12-1999
WO 9625183	A	22-08-1996	AU 4865796 A	04-09-1996
			WO 9625183 A2	22-08-1996
US 4704131	A	03-11-1987	JP 1473386 C	27-12-1988
			JP 58180162 A	21-10-1983
			JP 63020143 B	26-04-1988
			DE 3361391 D1	16-01-1986
			EP 0092414 A1	26-10-1983
WO 9606532	A	07-03-1996	AT 198119 T	15-01-2001
			AU 3341995 A	22-03-1996
			DE 69519675 D1	25-01-2001
			DE 69519675 T2	13-09-2001
			WO 9606532 A1	07-03-1996
			DK 778733 T3	02-04-2001
			EP 0778733 A1	18-06-1997

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Application No

PCT/US 01/04524

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9606532	A	ES 2154344 T3 JP 10505592 T PT 778733 T	01-04-2001 02-06-1998 29-06-2001
WO 9840091	A	17-09-1998	17-09-1998
US 5786324	A	28-07-1998	03-11-1998
JP 9010288	A	14-01-1997	NONE